



HANDBOOK

Partial Differential Equations of Applied Mathematics

Prepared by the Course Team

The Open University Press, Walton Hall, Milton Keynes

First published 1974. Reprinted 1975

Copyright © 1974 The Open University

All rights reserved. No part of this work may be reproduced in any form, by mimeograph or any other means, without permission in writing from the publishers.

Produced in Great Britain by
The Open University

ISBN 0 335 01255 8

This text forms part of the correspondence element of an Open University Third Level Course

For general availability of supporting material referred to in this text, please write to the Director of Marketing, The Open University, P.O. Box 81, Milton Keynes, MK7 6AT.

Further information on Open University courses may be obtained from The Admissions Office, The Open University, P.O. Box 48, Milton Keynes, MK7 6AB.

1.2

<u>Contents</u>	Page
1 Glossary	5
2 Vectors	36
3 Useful Formulas and Theorems	42
4 Table of Indefinite Integrals	50
5 Table of Definite Integrals	52

Note

Cross-references in this handbook are indicated by the use of CAPITALS. In addition, references are given to appearances in the course of definitions/formulas/theorems. The set books are denoted by S and W, and the correspondence text of Unit n is indicated by n.

1 GLOSSARY

ABSOLUTELY CONVERGENT The series $\sum_{n=1}^{\infty} a_n$ is absolutely convergent if
 6: page 13 the series $\sum_{n=1}^{\infty} |a_n|$ converges.
 14: page 7

AMPLIFICATION MATRIX The THREE-LEVEL SCHEME $Au_{j+1} = Bu_j + Cu_{j-1}$ is
 15: page 7 equivalent to a pair of two-level schemes
 which may be written in the form

$$\begin{bmatrix} u_{j+1} \\ v_{j+1} \end{bmatrix} = P \begin{bmatrix} u_j \\ v_j \end{bmatrix}$$

where

$$P = \begin{bmatrix} A^{-1}B & A^{-1}C \\ I & 0 \end{bmatrix}$$

P is called the amplification matrix.

ANGULAR FREQUENCY One-dimensional wave motion which is described
 7: page 12 by $A \sin(kx - \omega t)$, for example, where x represents
 the displacement and t the time, has angular
frequency ω , wave length $2\pi/k$ and wave number k.

ANGULAR MOMENTUM The angular momentum about the point P of a
 16: page 13 particle at the point Q with velocity \underline{v} and mass m is

$$m\underline{r} \times \underline{v}, \text{ where } \underline{r} = \underline{PQ}$$

ANGULAR VELOCITY The angular velocity about the point P of a
 16: page 12 particle at the point Q with velocity \underline{v} is
 $(\underline{r} \times \underline{v})/r^2$, where $\underline{r} = \underline{PQ}$; it measures the rate of
 rotation of the particle about Q.

ASYMPTOTIC RATE OF
CONVERGENCE

11: page 20

The asymptotic rate of convergence of the
ITERATIVE SCHEME

$$\tilde{x}^{(n+1)} = G\tilde{x}^{(n)} + Hb$$

is $-\log \rho$, where ρ is the SPECTRAL RADIUS of G .

BAND MATRIX

11: page 16

A band matrix is one in which all the nonzero
entries are confined to the leading diagonal and
those diagonals close to it. If a_{ij} is that
element of a matrix A which lies in row i and
column j , then A is a band matrix of band width
 $2k+1$ provided

$$a_{ij} = 0 \text{ for } |i-j| > k.$$

BAND WIDTH

See BAND MATRIX.

BESSEL'S EQUATION

14: page 6

W : page 179

Bessel's equation of order m is

$$t^2 \frac{d^2 u}{dt^2} + t \frac{du}{dt} + (t^2 - m^2)u = 0$$

or

$$\frac{d}{dt} \left(t \frac{du}{dt} \right) - \frac{m^2}{t} u + tu = 0.$$

BESSEL FUNCTION

W : page 179

A solution of BESSEL'S EQUATION (which is bounded
at the origin) is called a Bessel function (of the
first kind). A Bessel function of the first kind
of order m (>0) is given by

$$J_m(t) = \sum_{k=0}^{\infty} \frac{(-1)^k \left(\frac{1}{2}t\right)^{m+2k}}{k! \Gamma(m+k+1)}.$$

- BODY FORCE A body force is a force which acts on each element of a body from outside the body (e.g., gravitational force).
- 1 : page 10
- BOUNDARY CONDITIONS Boundary conditions are conditions at a spatial boundary for all times. For example, the equation
- 1: page 9
- $$\frac{\partial^2 u}{\partial x^2} = \frac{\partial^2 u}{\partial t^2} \quad x \in (\alpha, \beta), \quad t > 0,$$
- may be given with the boundary conditions
- $$u(\alpha, t) = f(t) \quad t \geq 0,$$
- $$\frac{\partial u}{\partial x}(\beta, t) = g(t) \quad t \geq 0.$$
- BOUNDARY VALUE PROBLEM See GREEN'S FUNCTION.
- CANONICAL FORM See STANDARD FORM.
- CAUCHY PROBLEM A Cauchy problem (or initial-boundary value problem) is a problem where the solution to a partial differential equation is required subject to the solution and its normal derivative both being specified on the initial line.
- 2: page 18
- 5: page 5
- CENTRAL-DIFFERENCE OPERATOR The central-difference operator δ_h is defined by
- $$\delta_h u : x \mapsto u(x + \frac{1}{2}h) - u(x - \frac{1}{2}h) \quad x \in \mathbb{R}.$$
- 5 : page 9
- CHARACTERISTICS See CLASSIFICATION OF OPERATORS.

CLASSIFICATION OF

The operator

OPERATORS

$$L:u \rightarrow A(x,t)\frac{\partial^2 u}{\partial t^2} + B(x,t)\frac{\partial^2 u}{\partial x \partial t} + C(x,t)\frac{\partial^2 u}{\partial x^2} + F\left(x,t,u,\frac{\partial u}{\partial t},\frac{\partial u}{\partial x}\right)$$

2: page 19W: page 43(a) hyperbolic at (\bar{x}, \bar{t}) if $[B(\bar{x}, \bar{t})]^2 - 4A(\bar{x}, \bar{t})C(\bar{x}, \bar{t}) > 0$,(b) parabolic at (\bar{x}, \bar{t}) if $[B(\bar{x}, \bar{t})]^2 - 4A(\bar{x}, \bar{t})C(\bar{x}, \bar{t}) = 0$,(c) elliptic at (\bar{x}, \bar{t}) if $[B(\bar{x}, \bar{t})]^2 - 4A(\bar{x}, \bar{t})C(\bar{x}, \bar{t}) < 0$.

For hyperbolic equations, the two families of curves which satisfy the equation

$$A\left(\frac{dx}{dt}\right)^2 - B\frac{dx}{dt} + C = 0$$

are called characteristics.

COEFFICIENT OF

See VISCOSITY.

VISCOSITY

COMPATIBLE

See CONSISTENT.

COMPLETE

See CONVERGENCE IN THE MEAN.

COMPLEX FOURIER
SERIES

The FOURIER SERIES expansion of a function (with domain R or C) in terms of the ORTHOGONAL BASIS

6: page 18

$$\{e^{inx} : n \in Z, x \in [-\pi, \pi]\}$$

is a complex Fourier series.CONDITIONALLY
STABLE

A FINITE-DIFFERENCE SCHEME is conditionally stable if it is STABLE under certain conditions (e.g., for particular values of the MESH RATIO) and unstable under others.

8: page 17

CONSISTENT

A FINITE-DIFFERENCE REPLACEMENT of a differential equation is consistent (or compatible) if its LOCAL TRUNCATION ERROR approaches zero as the MESH spacings tend to zero. (See also LAX'S THEOREM.)

8: page 9

CONSISTENT ORDERING

11: page 28

In the successive computation of a set of values $\{u_1, \dots, u_N\}$ using the GAUSS-SEIDEL or SOR iterative scheme, a scanning order of the points $1, 2, \dots, N$ is a consistent ordering if it yields the same set of equations (at each iteration) as would be obtained from some matrix which can be partitioned as

$$\begin{bmatrix} D_1 & F_1 & & & \\ E_1 & D_2 & F_2 & & \\ & & & & \\ & & E_{m-2} & D_{m-1} & F_{m-1} \\ & & & E_{m-1} & D_m \end{bmatrix} :$$

where the D_i are diagonal matrices and $m \geq 2$.

CONTINUOUS WITH
RESPECT TO DATAW : page 6

A problem is continuous with respect to its data if solutions corresponding to data which differ by small amounts also differ by small amounts.

CONTINUOUSLY
DIFFERENTIABLE

A function f is continuously differentiable on the interval I if its derived function f' is continuous on I .

CONVERGENCE

IN THE MEAN

W: page 71

The infinite sequence of functions s_1, s_2, \dots converges in the mean to f (on the interval $[a, b]$) with respect to the positive weight function ρ if

$$\lim_{N \rightarrow \infty} \int_a^b (f - s_N)^2 \rho = 0.$$

Let $\{\phi_n\}$ be an infinite sequence of functions ORTHOGONAL on $[a, b]$ with respect to the weight function ρ . For any function f let $s_N = \sum_{n=1}^N c_n \phi_n$ be the sum of the first N terms of the FOURIER SERIES of f . If the limit above is zero for every function f for which $\int_a^b f^2 \rho$ is finite, the set $\{\phi_n\}$ is said to be complete.

CONVERGENCE OF A
FINITE-DIFFERENCE
REPLACEMENT

8: page 8

A FINITE-DIFFERENCE REPLACEMENT of a differential equation is convergent if, as the MESH spacings tend to zero, the finite-difference solution tends to the true solution of the differential equation either at a fixed point, or for all points along the furthest time level under consideration.

CONVEX

3: page 14

A subset of R^2 (or R^3) is convex if no straight line intersects its boundary in more than two points, except possibly along a straight-line (or plane) section of the boundary.

CURVE

2: page 18W: page 49A curve C in R^2 is given by

$$\{(x,y): x = \phi(\tau), y = \theta(\tau) \quad \tau_0 \leq \tau \leq \tau_1\}$$

where ϕ and θ are continuous. If ϕ and θ are CONTINUOUSLY DIFFERENTIABLE on $[\tau_0, \tau_1]$ with $\theta'^2 + \phi'^2 > 0$ we say that C is a continuously differentiable curve. If these conditions hold except for a finite number of values of τ , C is a piecewise continuously differentiable curve. The curve C is closed if

$$\phi(\tau_1) = \phi(\tau_0) \text{ and } \theta(\tau_1) = \theta(\tau_0).$$

D'ALEMBERT'S SOLUTION

W: pages 9,13

D'Alembert's solution is the general solution of the one-dimensional wave equation

$$\frac{\partial^2 u}{\partial x^2} = \frac{1}{c^2} \frac{\partial^2 u}{\partial t^2}$$

in the form $u(x,t) = p(x+ct) + q(x-ct)$, where p and q are arbitrary functions.

DELTA FUNCTION

10: page 15The delta function in n dimensions is defined by

$$\delta(\underline{r}-\underline{\rho}) = 0 \quad \underline{r} \neq \underline{\rho}$$

$$\int_D f(\underline{r}) \delta(\underline{r}-\underline{\rho}) d\underline{r} = \begin{cases} f(\underline{\rho}) & \underline{\rho} \in D \\ 0 & \underline{\rho} \notin D \end{cases}$$

for all functions f and all domains $D \subset R^n$.

DIFFUSION EQUATION

3: page 31The diffusion equation is

$$\frac{\partial u}{\partial t} - k \nabla^2 u = 0.$$

DIRECTIONAL DERIVATIVE

3: page 7The directional derivative of the SCALAR FIELD

ϕ in the direction of the unit vector \underline{e} is given by the inner product $\underline{e} \cdot \underline{\text{grad}} \phi$.

- DIRICHLET PROBLEM A boundary value problem in which a solution, u ,
3: page 9 of a differential equation in a DOMAIN is
9: page 14 specified on the boundary C of the domain is called
 a Dirichlet problem. If, instead of u , the normal
 derivative $\partial u / \partial n$ is specified on C , the problem
 is called a Neumann problem.
- DISCRETIZATION Discretization is the replacement of a derivative
8: page 17 by a finite-difference formula (e.g., the FORWARD-
 DIFFERENCE FORMULA).
- DISPERSION The physical phenomenon of the velocity of a
7: page 13 wave being dependent upon its frequency is called
 dispersion.
- DISPLACEMENT VECTOR The displacement vector $\underline{\Delta}^{(n)}$ is the correction
11: page 20 to $\underline{x}^{(n)}$ at the $(n+1)$ th iteration of an ITERATIVE
5: page 78 SCHEME, i.e.,

$$\underline{\Delta}^{(n)} = \underline{x}^{(n+1)} - \underline{x}^{(n)}.$$
- DOMAIN (1) The domain of a given function f is ^{the} set of values
 for which f is defined.
- DOMAIN (2) A connected open subset of R^n is called a domain.
W: page 49 (A domain in one dimension is just an open interval.)
- DOMAIN OF DEPENDENCE The domain of dependence of the point (\bar{x}, \bar{t}) for a
2: page 12 given HYPERBOLIC equation is the set of all points
 in the solution domain whose DOMAINS OF INFLUENCE
 include the point (\bar{x}, \bar{t}) .

DOMAIN OF INFLUENCE	For a HYPERBOLIC equation, the two
$\tilde{2}$: page 12	CHARACTERISTIC curves through the point (\bar{x}, \bar{t})
\tilde{W} : page 40	bound a domain which is called the <u>domain of influence</u> of the point (\bar{x}, \bar{t}) . Values of the solution at \bar{x} for a given time \bar{t} can affect the solution only at points within this domain for which $t > \bar{t}$.
EIGENFUNCTION	See EIGENVALUE.
EIGENVALUE	Given a linear problem consisting of the
\tilde{W} : pages 65,160	equation $Lu + \lambda u = 0$ (where L is a LINEAR
\tilde{S} : page 62	OPERATOR) with homogeneous BOUNDARY CONDITIONS, it sometimes turns out that nontrivial solutions are possible only if λ takes on prescribed values called <u>eigenvalues</u> . For a given problem, the solutions associated with each eigenvalue are called <u>eigenfunctions</u> (in the case of a function space) or <u>eigenvectors</u> .
ELLIPTIC	See CLASSIFICATION OF OPERATORS.
EQUATION OF STATE	The <u>equation of state</u> of a fluid is a relation
$\tilde{1}$: page 15	between pressure and density. For an ideal gas, we have the adiabatic equation $p\rho^{-\gamma} = \text{constant}$.
EVEN FUNCTION	A function f with domain R or $[-a, a]$ or $(-a, a)$
\tilde{W} : page 23	is <u>even</u> if, for each x in the domain,
	$f(x) = f(-x).$

- EXPLICIT SCHEME A FINITE-DIFFERENCE SCHEME is explicit if each equation expresses one unknown PIVOTAL VALUE in terms of known pivotal values.
- S : page 11
- FINITE-DIFFERENCE REPLACEMENT A finite-difference replacement of a differential equation is a FINITE-DIFFERENCE SCHEME obtained by DISCRETIZATION of the derivatives in the equation.
- 5 : page 11
- FINITE-DIFFERENCE SCHEME (or METHOD) A finite-difference scheme is a formula relating the PIVOTAL VALUES of a function over a MESH.
- 5 : page 11
- FINITE FOURIER TRANSFORM The finite Fourier transform of $u(r, \theta)$, where $-\pi \leq \theta < \pi$, is the sequence $\{a_n(r), b_n(r)\}$ given by
- $$a_n(r) = \frac{1}{\pi} \int_{-\pi}^{\pi} u(r, \theta) \cos n\theta \, d\theta \quad n = 0, 1, 2, \dots$$
- $$b_n(r) = \frac{1}{\pi} \int_{-\pi}^{\pi} u(r, \theta) \sin n\theta \, d\theta \quad n = 1, 2, \dots$$
- W : page 129
- FINITE SINE TRANSFORM The finite sine transform of $u(r, \theta)$, where $0 < \theta < \pi$, is the sequence $\{b_n(r)\}$ given by
- $$b_n(r) = \frac{2}{\pi} \int_0^{\pi} u(r, \theta) \sin n\theta \, d\theta \quad n = 1, 2, \dots$$
- W : page 127
- FLUID EQUATIONS See EQUATION OF STATE, MASS-CONSERVATION EQUATION and MOMENTUM EQUATION.

FLUX

3: page 2116: page 22

The flux of a fluid is its rate of flow, i.e., the volume crossing a given surface per unit time. If the velocity of the fluid is given by the VECTOR FIELD \underline{v} , the flux across the surface S is given by

$$\int_S \underline{v} \cdot \underline{n} \, dS$$

where \underline{n} is a unit vector normal to S . In the case of a fluid which flows down a pipe with speed w the flux through a cross-section D of the pipe is

$$\int_D w \, dA.$$

FORWARD-DIFFERENCE
OPERATOR

5: page 9

The forward-difference operator Δ_h is defined by

$$\Delta_h u : x \mapsto u(x+h) - u(x) \quad x \in \mathbb{R}.$$

FOURIER-BESSEL SERIES

14: page 10

A Fourier-Bessel series is the FOURIER SERIES expansion of an arbitrary function with domain $(0,1)$ as $\sum_{k=1}^{\infty} c_k J_m(\sqrt{\lambda_k^{(m)}} x)$ in terms of BESSEL FUNCTIONS satisfying $J_m(\sqrt{\lambda_k^{(m)}}) = 0$.

FOURIER COEFFICIENT

See FOURIER SERIES.

FOURIER SERIES

W: page 72

The expansion of an arbitrary SQUARE INTEGRABLE function f as a series $\sum_{k=1}^{\infty} c_k \phi_k$ in terms of the ORTHOGONAL set of functions $\{\phi_k\}$, with

$$c_k = \frac{f \cdot \phi_k}{\phi_k \cdot \phi_k},$$

is called a Fourier series with Fourier coefficients $\{c_k\}$.

GAMME FUNCTION

The Gamma function is the function

14: page 7

$$\Gamma : a \mapsto \int_0^{\infty} e^{-x} x^{a-1} dx \quad a > 0$$

GAUSS-SEIDEL

For the ITERATIVE SCHEME

ITERATION MATRIX

$$\underline{x}^{(n+1)} = L\underline{x}^{(n+1)} + U\underline{x}^{(n)} + \underline{b},$$

11: page 23

the matrix $(I-L)^{-1}U$ is called the Gauss-Seidel iteration matrix.

S: page 78

GAUSS-SEIDEL

The Gauss-Seidel method for solving the problem

METHOD

$$(I-L-U)\underline{x} = \underline{b},$$

11: pages 23,24

where L and U are respectively lower and upper triangular matrices, is given by the ITERATIVE SCHEME

S: page 78

$$\underline{x}^{(n+1)} = L\underline{x}^{(n+1)} + U\underline{x}^{(n)} + \underline{b}.$$

GLOBAL ERROR

The global error, ϵ_m at a MESH point, x_m a FINITE-DIFFERENCE REPLACEMENT of a differential equation is the difference between the computed finite-difference solution and the true solution to the differential equation, at that point.

S: page 27

8: page 7

GREEN'S FUNCTION FOR
ORDINARY DIFFERENTIAL
EQUATIONS

W : page 122

10 : page 16

Green's function $G(x, \xi)$ for the boundary value
problem

$$\frac{d}{dx} \left[p(x) \frac{du}{dx} \right] + q(x)u(x) = -f(x) \quad x \in (\alpha, \beta)$$

$$u(\alpha) = u(\beta) = 0$$

is given, for each $\xi \in (\alpha, \beta)$, by the continuous
solution of

$$\frac{d}{dx} \left[p(x) \frac{dG}{dx} \right] + q(x)G = 0 \quad x \neq \xi,$$

$$G \Big|_{x=\alpha} = G \Big|_{x=\beta} = 0,$$

$$G \Big|_{x=\xi+0} - G \Big|_{x=\xi-0} = 0,$$

$$\frac{dG}{dx} \Big|_{x=\xi+0} - \frac{dG}{dx} \Big|_{x=\xi-0} = -\frac{1}{p(\xi)}.$$

Alternatively, we can write this system as

$$\frac{d}{dx} \left[p(x) \frac{dG}{dx} \right] + q(x)G = -\delta(x-\xi),$$

$$G(\alpha, \xi) = G(\beta, \xi) = 0,$$

where $\delta(x-\xi)$ is the DELTA FUNCTION in one
dimension. The solution to the boundary value
problem above may be expressed as

$$u(x) = \int_{\alpha}^{\beta} G(x, \xi) f(\xi) d\xi.$$

GREEN'S FUNCTION

For any DOMAIN D with boundary C , Green's function

FOR PARTIAL

$G(\underline{r}; \underline{\rho})$ for the boundary value problem

DIFFERENTIAL

$$L[u](\underline{r}) = -F(\underline{r}) \quad \underline{r} \in D$$

EQUATIONS

$$u(\underline{r}) = 0, \quad \underline{r} \in C$$

\underline{W} : page 135

where L is a linear ELLIPTIC operator, is given,

$\underline{10}$: page 18

for each $\underline{\rho} \in D$, by the continuous solution of

$$L[G](\underline{r}; \underline{\rho}) = -\delta(\underline{r} - \underline{\rho}) \quad \underline{r} \in D$$

$$G(\underline{r}; \underline{\rho}) = 0 \quad \underline{r} \in C.$$

The solution to the problem above may be written

in the form

$$u(\underline{r}) = \int_D G(\underline{r}; \underline{\rho}) F(\underline{\rho}) d\underline{\rho}.$$

GRID

See MESH.

HARMONIC

The n th harmonic of a function f with domain

$\underline{6}$: page 14

$[-L, L]$ is $a_n \sin n\pi x/L + b_n \cos n\pi x/L$, where a_n and b_n

$\underline{16}$: page 20

are the appropriate FOURIER COEFFICIENTS of f .

HARMONIC FUNCTION

A solution of Laplace's equation

\underline{W} : page 52

$$\nabla^2 u = 0$$

is called a harmonic function.

HEAT CONTENT

Let $u(x, y, z, t)$ be the temperature at the point

$\underline{3}$: page 31

(x, y, z) in the domain D at time $t \geq 0$. The heat

content of D is a measure of the total amount of

heat in D and is given by

$$c\rho \iiint_D u(x, y, z, t) \, dx dy dz$$

where ρ is the density and c the SPECIFIC HEAT

of the material.

HEAT EQUATION

W: page 58

The heat equation is another name for the
DIFFUSION EQUATION.

HÖLDER CONTINUOUS

W: page 79

The function $f : I \rightarrow \mathbb{R}$ is Hölder continuous
at $x \in I$ if $\exists M > 0, \alpha > 0$ such that $\forall y \in I$

$$|f(y) - f(x)| \leq M|y-x|^\alpha.$$

HOMOGENEOUS EQUATION

W: page 30

If L is any LINEAR operator then

$$Lx = 0$$

is a homogeneous equation.

HOOKE'S LAW

W: page 6

Hooke's Law states that force exerted by an
elongated elastic medium is proportional to
the extension per unit length; the constant of
proportionality is called Young's Modulus.

HYPERBOLIC

See CLASSIFICATION OF OPERATORS.

IDEAL FLUID

16: page 7

An ideal fluid is one whose coefficient of
VISCOSITY is zero.

IMPLICIT SCHEME

S: page 18

A FINITE-DIFFERENCE SCHEME is implicit if it
must be solved for several unknown values
simultaneously.

IMPROPER INTEGRAL

6: page 14

A integral in which the integrand or the domain
of integration is unbounded may be evaluated as
an improper integral if it may be realized as
the limit of a sequence of (proper) integrals.
For example, if $f(a+0)$ does not exist we may
define

$$\int_a^b f = \lim_{\epsilon \rightarrow 0^+} \int_{a+\epsilon}^b f$$

provided this limit exists. Similarly we define

$$\int_a^\infty f = \lim_{b \rightarrow \infty} \int_a^b f$$

provided this limit exists

INDUCED INSTABILITY A method of solution for a problem suffers from
 5 : page 6 induced instability if the method magnifies LOCAL
 ERRORS so that the computed result differs
 significantly from the true result.

INFLUENCE FUNCTION The influence function $R(x, \xi)$ for the initial value
 W : page 119 problem

$$\frac{d}{dx} \left[p(x) \frac{du}{dx} \right] + q(x)u(x) = f(x) \quad x > \alpha,$$

$$u(\alpha) = u'(\alpha) = 0,$$

is given, for each $\xi > \alpha$, by the solution of the
 initial value problem

$$\frac{d}{dx} \left[p(x) \frac{dR}{dx} \right] + q(x)R = 0 \quad x > \xi,$$

$$R|_{x=\xi} = 0,$$

$$\frac{dR}{dx} \Big|_{x=\xi} = \frac{1}{p(\xi)},$$

or the equivalent problem

$$\frac{d}{dx} \left[p(x) \frac{dR}{dx} \right] + q(x)R = \delta(x-\xi),$$

$$R|_{x=\xi} = 0.$$

The solution of the original problem is given by

$$u(x) = \int_{\alpha}^x R(x, \xi) f(\xi) d\xi.$$

INHERENT INSTABILITY	A mathematical problem suffers from <u>inherent instability</u> if it is not CONTINUOUS WITH RESPECT TO ITS DATA or is not WELL CONDITIONED.
<u>5</u> : page 5	
INITIAL-BOUNDARY VALUE PROBLEM	See CAUCHY PROBLEM.
INITIAL CONDITIONS	<u>Initial conditions</u> for a partial differential equation are conditions given at spatial points for time $t = t_0$. For example, a stretched string may have its motion started with a given shape, $u(x,0) = f(x)$, and a given velocity distribution $\frac{\partial u}{\partial t}(x,0) = g(x)$. For ordinary differential equations, <u>initial conditions</u> are conditions specified at one point of the solution domain.
<u>1</u> : page 9	
INITIAL VALUE PROBLEM	See INFLUENCE FUNCTION.
ITERATIVE SCHEME	An <u>iterative scheme</u> for solving a problem is one in which the solution is obtained by a process of successive approximation.
<u>11</u> : page 15	
JACOBIAN	The Jacobian of the N functions f_1, f_2, \dots, f_N each with domain variable $(x_1, x_2, \dots, x_N) \in \mathbb{R}^N$ and codomain \mathbb{R} is the determinant of the matrix
<u>15</u> : page 15	

$$\begin{bmatrix} \frac{\partial f_1}{\partial x_1} & \dots & \frac{\partial f_1}{\partial x_N} \\ \vdots & & \vdots \\ \frac{\partial f_N}{\partial x_1} & \dots & \frac{\partial f_N}{\partial x_N} \end{bmatrix}$$

JACOBI METHOD

11: page 245: page 78The Jacobi method is the ITERATIVE SCHEME

$$\underline{x}^{(n+)} = L\underline{x}^{(n)} + U\underline{x}^{(n)} + \underline{b}$$

for the solution of the equation

$$(I-L-U)\underline{x} = \underline{b},$$

where L and U are respectively lower and upper triangular matrices.

JUMP DISCONTINUITY

6: page 14

The function f has a jump discontinuity at x if the one-sided limits $f(x+0)$ and $f(x-0)$ exist but are not both equal to $f(x)$.

KINETIC ENERGY

2: page 14

The kinetic energy of a particle is the energy it possesses by virtue of its motion, and is defined to be

$$\frac{1}{2}(\text{mass}) \times (\text{velocity})^2.$$

KRONECKER DELTA

The Kronecker delta is given by

$$\delta_{ij} = \begin{cases} 0 & i \neq j \\ 1 & i = j \end{cases} \quad i, j \in \mathbb{Z}.$$

LAPLACE'S EQUATION

W: pages 43,49,52Laplace's equation is $\nabla^2 u = 0$.LAW OF CONSERVATION
OF ENERGY2: page 13

The Law of Conservation of Energy states that the change in the total energy (= KINETIC ENERGY + POTENTIAL ENERGY) of a system is equal to the work done on the system by all the forces. In particular the energy of an isolated system remains constant.

- LAW OF CONSERVATION OF MASS
1: page 12
 The Law of Conservation of Mass states that the total mass of a quantity of material remains constant in time, however it becomes distributed in space.
- LINE INTEGRAL
3: page 12
 The line integral $\int_C f \, ds$ represents $\lim_{\Delta s \rightarrow 0} \sum f_P \Delta s$ over the CURVE C, where f_P is the value of f at a point P in the small element of C with length Δs .
- LINEAR (OPERATOR)
W: page 29
 L is a linear transformation (operator) if it has the property
- $$L[\alpha \underline{u} + \beta \underline{v}] = \alpha L[\underline{u}] + \beta L[\underline{v}],$$
- where α and β are any constants and $\underline{u}, \underline{v}$ are any vectors (functions).
- LINEAR PARTIAL DIFFERENTIAL EQUATION
W: page 30
 $L[\underline{u}] = F$
 where L is a LINEAR OPERATOR involving partial differentiation and F is a given function.
- LOCAL ERROR
5: page 27
 The local error is the error made at a point when a FINITE-DIFFERENCE SCHEME is used once only. In the complete step-by-step process, the finite-difference scheme is used many times so that local errors accumulate to produce GLOBAL ERRORS.
- LOCAL TRUNCATION ERROR
5: page 14
 The local truncation error (at a point) in a FINITE-DIFFERENCE REPLACEMENT of a differential equation is the LOCAL ERROR introduced by DISCRETIZATION.

MASS-CONSERVATION
EQUATION

1: page 13

14: page 26

The Mass-Conservation equation expresses mathematically the physical LAW OF CONSERVATION OF MASS as applied to problems in fluid flow. It is given by

$$\frac{\partial \rho}{\partial t} + \operatorname{div} \rho \underline{u} = 0,$$

where ρ and \underline{u} denote the density and velocity respectively, and t is the time coordinate. In one spatial dimension with coordinate x the equation reduces to

$$\frac{\partial \rho}{\partial t} + \rho \frac{\partial u}{\partial x} + u \frac{\partial \rho}{\partial x} = 0.$$

MEAN SQUARE
DEVIATION

W: page 71

The mean square deviation of g from f , with respect to the weight function ρ on $[a,b]$, is

$$\int_a^b (f-g)^2 \rho.$$

MESH

5: page 7

The set of points $\{(x_i, t_j)\} = \{(x_0 + ih, t_0 + jk)\}$ where h and k are constants is called a mesh or grid. In a given mesh each point (x_i, t_j) is called a mesh point, and h and k are called the mesh spacings or mesh lengths in the x - and t -directions respectively.

MESH RATIO

5: pages 11,25

A FINITE-DIFFERENCE REPLACEMENT of a HYPERBOLIC or PARABOLIC differential equation depends on the MESH spacings h and k (in the x - and t -directions respectively) according to the value of the mesh ratio k^2/h^2 (for hyperbolic equations) or k/h^2 (for parabolic equations). For typical examples, see Section 3.

MOLECULAR DIAGRAM

5: page 125: page 11A molecular diagram is a diagrammatic

representation of a FINITE-DIFFERENCE SCHEME

which indicates the coefficients of the PIVOTAL VALUES related by the scheme.

MOMENT

16: page 13The moment about the point P of a force, actingthrough the point Q and represented by the geometric vector \underline{F} , is $\underline{r} \times \underline{F}$ where \underline{r} is the geometric vector \underline{PQ} .

MOMENTUM EQUATION

1: page 1514: page 27The Momentum equation expresses mathematically NEWTON'S SECOND LAW OF MOTION as applied to fluid flow. It is given by

$$\frac{\partial \underline{u}}{\partial t} + (\underline{u} \cdot \underline{\text{grad}}) \underline{u} + \frac{1}{\rho} \underline{\text{grad}} p = \underline{F},$$

where ρ, \underline{u}, p and \underline{F} denote the density, velocity, pressure and BODY FORCE per unit mass respectively, and t is the time coordinate. In one spatial dimension with coordinate x the equation reduces to

$$\frac{\partial u}{\partial t} + u \frac{\partial u}{\partial x} + \frac{1}{\rho} \frac{\partial p}{\partial x} = F.$$

NEUMANN PROBLEM

See DIRICHLET PROBLEM.

NEWTON

A newton is the SI unit of force.

NEWTONIAN FLUID

16: page 7A Newtonian fluid is one whose coefficient of VISCOSITY does not depend on the velocity gradient.

- ORTHONORMAL FUNCTIONS A set of functions $\{\phi_n\}$ which satisfies
- 6 : page 12
$$\int_{\alpha}^{\beta} \phi_m(x) \phi_n(x) \rho(x) dx = \delta_{mn},$$
- where δ_{mn} is the KRONECKER DELTA, is said to be orthonormal with respect to the weight function ρ on the interval (α, β) .
- OVER-RELAXATION See SUCCESSIVE OVER-RELAXATION METHOD.
- FACTOR
- PARABOLIC See CLASSIFICATION OF EQUATIONS.
- PERIODIC FUNCTION A function f is periodic (with period a) if, for all $x \in \mathbb{R}$, $f(x) = f(x+a)$.
- PIECEWISE CONTINUOUS A function is piecewise continuous on $[a, b]$ if it is continuous at each point of $[a, b]$ except for a finite number of JUMP DISCONTINUITIES.
- 6 : page 15
- PIECEWISE CONTINUOUSLY See CURVE.
- DIFFERENTIABLE CURVE
- PIVOTAL VALUE The values of a function u at MESH points are called pivotal values of u .
- 5 : page 7
- POINTWISE CONVERGENCE Given an infinite sequence of functions ϕ_1, ϕ_2, \dots with domain I , we let
- (OF A SERIES) $s_N(x) = \sum_{n=1}^N c_n \phi_n(x)$. If for each $x \in I$
- W : page 70 $\lim_{N \rightarrow \infty} s_N(x) = f(x)$, we say that the series $\sum_{n=1}^{\infty} c_n \phi_n(x)$ converges to $f(x)$ pointwise in I .

POISE A poise is the cgs unit of VISCOSITY.

16: page 6

POISSON'S EQUATION Poisson's equation is

W : pages 49,50,129

$$\nabla^2 u = -F.$$

3 : page 5

POTENTIAL ENERGY The potential energy of a system is the energy

2 : page 14

stored within the system.

PROPERLY POSED

A system consisting of a partial differential

PROBLEM

equation together with (INITIAL and) BOUNDARY

W : page 6

CONDITIONS is properly posed if its solution

3 : page 10

(a) exists, (b) is unique and (c) is CONTINUOUS WITH RESPECT TO THE DATA.

PROPERTY A

A square matrix is said to have Property A if,

11: page 29

by transposing pairs of rows and pairs of corresponding columns, it can be transformed to the form of the matrix

$$\begin{bmatrix} D_1 & F \\ E & D_2 \end{bmatrix}$$

where D_1 and D_2 are diagonal matrices.

RAYLEIGH QUOTIENT

Consider the differential equation

W : pages 163,165

$$(pu')' - qu + \lambda pu = 0 \quad \text{in } (\alpha, \beta),$$

where q and p are continuous and p is CONTINUOUSLY DIFFERENTIABLE in (α, β) , along with $u(\alpha) = u(\beta) = 0$.

Let ϕ vanish at α and β and be twice continuously differentiable in (α, β) . The Rayleigh quotient of ϕ is given by the ratio

$$\frac{\int_{\alpha}^{\beta} (p\phi'^2 + q\phi^2)}{\int_{\alpha}^{\beta} \phi^2}$$

Characterizations of the EIGENVALUES of the problem above as minima of the Rayleigh quotient are called minimum principles for the eigenvalues.

RECURRENCE RELATION

A recurrence relation is an equation relating successive members of a sequence.

ROUNDING ERROR

A rounding error is introduced into the solution of a problem by working to a definite number of significant figures.

RESIDUAL VECTOR

11: page 16

$$A\underline{x} = \underline{b},$$

S: pages 31,79

then, if $\bar{\underline{x}}$ is an approximation to \underline{x} , the vector

$$\underline{r} = \underline{b} - A\bar{\underline{x}}$$

is called the residual vector.

SCALAR FIELD

3: page 6

Let D denote a subset of R^3 ; then a function

$$\begin{array}{ccc} \phi : D \rightarrow R & & \phi : D \times R_0^+ \rightarrow R \\ & \text{or} & \\ \phi : (x,y,z) \mapsto u & & \phi : (x,y,z,t) \mapsto u \end{array}$$

is called a scalar field. (In the second case we say that ϕ is time-dependent.)

SELF-ADJOINT FORM

W: page 117

An ordinary differential equation written in the form

$$(pu')' + qu = f$$

on the interval (a,b) is said to be in self-adjoint form. We require that p be CONTINUOUSLY DIFFERENTIABLE and positive and that q and f be continuous on $[a,b]$.

SINGULAR POINT A singular point of the ordinary differential
10: page 7 equation

$$a(x)\frac{d^2u}{dx^2} + b(x)\frac{du}{dx} + c(x)u = 0$$

is a point x where either $a(x) = 0$ or $b(x)$ or $c(x)$ is infinite.

SOR See SUCCESSIVE OVER-RELAXATION METHOD.

SPARSE A matrix is sparse if it has a large proportion of
11: page 15 zero elements.

SPECIFIC HEAT The specific heat of a material is the quantity of
3: page 31 heat required to raise the temperature of unit mass of the material by one unit.

SPECTRAL RADIUS The spectral radius of the matrix M is

8: page 18

$$\rho(M) = \max_i |\lambda_i|$$

5: page 79

where $\lambda_i (i=1,2,\dots,n)$ are the EIGENVALUES of M .

SQUARE INTEGRABLE The function f is square integrable on (α,β) with
6: page 13 respect to the weight function ρ if the integral

$$\int_{\alpha}^{\beta} f^2 \rho$$

exists.

STABLE A FINITE-DIFFERENCE SCHEME is stable if LOCAL ERRORS
8: page 17 do not accumulate exponentially as the step-by-step solution progresses, i.e., if it does not suffer from INDUCED INSTABILITY.

STANDARD FORM

W: pages 42,43

The standard form (or canonical form) for a

linear partial differential operator which is

(a) HYPERBOLIC is $\frac{\partial^2 u}{\partial \xi \partial \eta} + F\left(\frac{\partial u}{\partial \xi}, \frac{\partial u}{\partial \eta}, u, \xi, \eta\right) = 0;$

(b) PARABOLIC is $\frac{\partial^2 u}{\partial \eta^2} + F\left(\frac{\partial u}{\partial \xi}, \frac{\partial u}{\partial \eta}, u, \xi, \eta\right) = 0;$

(c) ELLIPTIC is $\frac{\partial^2 u}{\partial \xi^2} + \frac{\partial^2 u}{\partial \eta^2} + F\left(\frac{\partial u}{\partial \xi}, \frac{\partial u}{\partial \eta}, u, \xi, \eta\right) = 0.$

STANDING WAVE

1 : page 35

2 : page 9

A standing wave is a wave which does not move

laterally, but merely changes its amplitude

with time. A standing wave is easily recognized

by observing points of zero displacement which

remain stationary for all time, e.g., $A \sin \omega t \cos kx$.

SUBSIDIARY CONDITIONS

Subsidiary conditions are INITIAL and/or

BOUNDARY CONDITIONS.

SUCCESSIVE OVER- RELAXATION METHOD

(SOR)

11: page 26

The successive over-relaxation method for solving

$$(I - L - U)\underline{x} = \underline{b},$$

where L and U are respectively lower and upper triangular matrices, is given by the ITERATIVE SCHEME

$$(I - \omega L)\underline{x}^{(n+1)} = \{(1 - \omega)I + \omega U\}\underline{x}^{(n)} + \omega \underline{b}.$$

The parameter ω is called the over-relaxation factor, and determines the ASYMPTOTIC RATE

OF CONVERGENCE of the method. The optimum

over-relaxation factor is that value which

maximizes the rate of convergence of the method.

- SURFACE INTEGRAL** The surface integral $\int_S f \, dA$ of the function f over the surface S represents $\lim_{\Delta A \rightarrow 0} \sum f_P \Delta A$, where f_P is the value of f at a point P in the small element of S with area ΔA .
- 3:** pages 11,27
- TAYLOR APPROXIMATION** A Taylor approximation of a function is obtained by neglecting the remainder term in TAYLOR'S THEOREM (see Section 3). The first-order Taylor approximations for functions of one, two and three variables are respectively:
- 1:** page 13
- 2:** page 21
- 14:** page 26

$$u(x+h) \approx u(x) + hu'(x);$$

$$u(x+h, t+h) \approx \left[u + h \frac{\partial u}{\partial x} + k \frac{\partial u}{\partial t} \right]_{(x,t)};$$

$$u(x+\Delta x, y+\Delta y, z+\Delta z) \approx [u + \Delta \underline{r} \cdot \underline{\text{grad}} \, u]_{(x,y,z)}$$

$$\text{where } \Delta \underline{r} = (\Delta x, \Delta y, \Delta z).$$

- THERMAL CONDUCTIVITY** The thermal conductivity of a material is the rate at which heat flows across a temperature gradient of unit magnitude.
- W:** page 58

- THREE-LEVEL SCHEME** A three-level FINITE-DIFFERENCE SCHEME involves terms along the $(j-1)$ th, j th and $(j+1)$ th time levels. If it is linear, it may be written in matrix form as
- 15:** page 6

$$A \underline{u}_{j+1} = B \underline{u}_j + C \underline{u}_{j-1}.$$

TRAVELLING WAVE

1: page 20

A travelling wave is a wave which moves in a given direction without changing its shape. For example $f(x+ct)$ represents a wave travelling in the negative x -direction. The substitution $X = x+ct$ gives $f(X)$ which may represent a wave of fixed shape. Since the origin of the X -coordinate is at $x = -ct$, the wave travels with velocity c in the negative x -direction. Similarly, $f(x-ct)$ represents a travelling wave of velocity c in the positive x -direction.

TRIDIAGONAL MATRIX

5: page 18

A tridiagonal matrix is one whose nonzero elements appear on and adjacent to its main diagonal, i.e., a BAND MATRIX of BAND WIDTH 3.

UNIFORMLY BOUNDED

6: page 21

The infinite sequence of functions $\{b_n\}$ is uniformly bounded on the interval I if $\exists c \in \mathbb{R}$, independent of $n \in \mathbb{Z}^+$ and $x \in I$, such that $n \in \mathbb{Z}^+$ and $x \in I$, $|b_n(x)| < c$.

UNIFORM CONVERGENCE

2: page 106: page 9W: page 70

The infinite sequence of functions $\{f_n\}$ converges uniformly to f on the interval I if it is convergent to f in the uniform norm, i.e., if for each $\varepsilon > 0$ $\exists N_\varepsilon \in \mathbb{Z}^+$ such that

$$\max_{x \in I} |f(x) - f_N(x)| < \varepsilon$$

whenever $N \geq N_\varepsilon$. An equivalent formulation is: for each $\varepsilon > 0$, $\exists N_\varepsilon \in \mathbb{Z}^+$ independent of x such that $|f(x) - f_N(x)| < \varepsilon$ for all $x \in I$ whenever $N \geq N_\varepsilon$.

The infinite series of functions $\sum_{n=1}^{\infty} c_n \phi_n$ converges uniformly on the interval I if the sequence $\{s_N\}$ of partial sums

$$s_N = \sum_{n=1}^N c_n \phi_n$$

is uniformly convergent.

VECTOR FIELD

3: page 6

A vector field in two (or three) dimensions is a function with domain D in \mathbb{R}^2 (or \mathbb{R}^3) and codomain the space G^2 (or G^3) of geometric vectors. A time-dependent vector field is defined similarly with domain $D \times \mathbb{R}_0^+$.

VELOCITY POTENTIAL

14: page 15

If the velocity \underline{v} of a fluid is related to a SCALAR FIELD Φ by $\underline{v} = -\underline{\text{grad}}\Phi$, Φ is called the velocity potential of the flow, which is then said to be irrotational.

VISCOSITY

3: page 20

16: page 6

If a relative motion occurs in a fluid, a measurable resistance is experienced, and the fluid is said to exhibit viscosity or internal friction. For a fluid moving in parallel layers, the frictional force F on a plane surface of area A parallel to the fluid flow is given to a good approximation by

$$\frac{F}{A} = \mu \frac{\partial v}{\partial z}$$

where $\frac{\partial v}{\partial z}$ is the velocity gradient perpendicular to the direction of motion and μ is called the coefficient of viscosity. If ρ is the density of the fluid, then its kinematic viscosity is μ/ρ .

WAVE EQUATION

The wave equation isW : pages 9,36,48,66,152

$$\frac{\partial^2 u}{\partial t^2} = c^2 \nabla^2 u.$$

l : page 8

WAVE LENGTH

See ANGULAR FREQUENCY.

WAVE NUMBER

See ANGULAR FREQUENCY.

WELL CONDITIONED

The linear equation

ll: page 15

$$A\underline{u} = \underline{b}$$

is well conditioned if small changes in the elements of A and \underline{b} do not cause large changes in the solution vector \underline{u} .

YOUNG'S MODULUS

See HOOKE'S LAW.

2 VECTORS

A **vector space** V over a *field* F of scalars is a set of elements with an internal binary operation $+: V \times V \rightarrow V$ and an external binary operation of multiplication $F \times V \rightarrow V$ satisfying the following axioms for all $a, b, c \in V$ and $m, n \in F$:

- 1 $(a + b) + c = a + (b + c)$
- 2 $a + b = b + a$
- 3 $\exists 0 \in V$ such that $0 + a = a \quad \forall a \in V$
- 4 $\exists -a \in V$ such that $a + (-a) = 0$
- 5 $m(a + b) = ma + mb$
- 6 $(m + n)a = ma + na$
- 7 $(mn)a = m(na)$
- 8 $1a = a$

When $F = R$ (the reals) we talk about a *real vector space*; when $F = C$ (the complex numbers) we have a *complex vector space*.

An important notion in a vector space is that of linear independence. A set $\{v_1, \dots, v_n\}$ of vectors in V is **linearly independent** if


$$x_1 v_1 + x_2 v_2 + \dots + x_n v_n = 0 \quad (x_i \in F)$$

holds only for $x_1 = x_2 = \dots = x_n = 0$. Suppose now that $\{v_1, \dots, v_m\}$ is a *maximal* linearly independent set in V , i.e. if another vector in V is included in the set we obtain a set which is linearly dependent. Then $\{v_1, \dots, v_m\}$ is called a **basis** for V , and every other basis for V has precisely m vectors. We say that V has **dimension** m . It can also be shown that an arbitrary element $v \in V$ can be expressed uniquely in the form

$$v = x_1 v_1 + \dots + x_n v_n \quad (x_i \in F).$$

Geometric vectors are equivalence classes of parallel arrows of equal length. If \overrightarrow{AB} is an arrow, we write $\underline{\overrightarrow{AB}}$ for the geometric vector to which it belongs. The sum of two geometric vectors is obtained by choosing suitable arrows which can be added according to the *triangle law of addition*:

$$\underline{\overrightarrow{OA}} + \underline{\overrightarrow{AB}} = \underline{\overrightarrow{OB}}.$$



If λ is a real number then we define the product $\lambda \underline{\overrightarrow{AB}}$ to be the geometric vector whose arrows are parallel to those of $\underline{\overrightarrow{AB}}$ but with lengths multiplied by λ .

With respect to these definitions of addition and scalar multiplication the set G^3 of geometric vectors in space forms a real vector space. We often consider the set G^2 of geometric vectors in the plane, which is a *subspace* of this vector space. The null vector 0 is the geometric vector whose arrows are all of zero length.

Let Ox_1x_2 be a frame of rectangular Cartesian coordinate axes and let i, j, k be geometric vectors whose arrows are of unit length and parallel to the coordinate axes Ox_1, Ox_2, Oz respectively. Then every geometric vector in the plane can be written uniquely in the form

$$a = a_1 i + a_2 j,$$

and every geometric vector in space can be written uniquely in the form

$$a = a_1 i + a_2 j + a_3 k.$$

Thus $\{i, j\}$ forms a basis for the space G^2 of geometric vectors in the plane, and $\{i, j, k\}$ forms a basis for G^3 ; a_1, a_2, a_3 are, respectively, the x -, y -, z -coordinates of a .

If I is an interval on the real line or a domain in \mathbb{R}^2 or \mathbb{R}^3 , then the set of all functions $I \rightarrow \mathbb{R}$ is a real vector space with addition and scalar multiplication of functions given by

$$(f+g) : x \mapsto f(x) + g(x) \quad x \in I,$$

$$\forall \lambda \in \mathbb{R} \quad (\lambda f) : x \mapsto \lambda[f(x)] \quad x \in I.$$

Replacing \mathbb{R} by \mathbb{C} in the above definition yields a complex vector space of functions.

The space of all VECTOR FIELDS with domain D forms a real vector space with addition and scalar multiplication of fields given by

$$(\underline{u}+\underline{v}) : \underline{x} \mapsto \underline{u}(\underline{x}) + \underline{v}(\underline{x}) \quad \underline{x} \in D,$$

$$\forall \lambda \in \mathbb{R} \quad \lambda \underline{v} : \underline{x} \mapsto \lambda[\underline{v}(\underline{x})] \quad \underline{x} \in D.$$

An important property of a geometric vector is its *length* or *magnitude*. To define this more generally, we introduce the inner product.

A mapping $\cdot : V \times V \rightarrow \mathbb{R}$, where V is a real vector space, is called a (real) **inner product** if for all $a, b, c \in V$ and $\lambda, \mu \in \mathbb{R}$:

$$1 \quad a \cdot a \geq 0 \Leftrightarrow a = 0;$$

$$2 \quad a \cdot a \geq 0;$$

$$3 \quad a \cdot b = b \cdot a;$$

$$4 \quad a \cdot (\lambda b + \mu c) = \lambda a \cdot b + \mu a \cdot c.$$

We may specify an inner product on G^2 or G^3 by

$$\overrightarrow{OA} \cdot \overrightarrow{OB} = |\overrightarrow{OA}| |\overrightarrow{OB}| \cos \theta,$$

where $|\overrightarrow{OA}|, |\overrightarrow{OB}|$ denotes the length of the arrow $\overrightarrow{OA}, \overrightarrow{OB}$ respectively and θ is the angle between \overrightarrow{OA} and \overrightarrow{OB} . The geometric vectors i, j, k are *orthogonal*, i.e.

$$i \cdot j = j \cdot k = k \cdot i = 0.$$

(In fact, any two geometric vectors which are represented by perpendicular arrows are orthogonal.) If $\overrightarrow{OA} = a_1 \underline{i} + a_2 \underline{j} + a_3 \underline{k}$ and $\overrightarrow{OB} = b_1 \underline{i} + b_2 \underline{j} + b_3 \underline{k}$, then

$$\overrightarrow{OA} \cdot \overrightarrow{OB} = a_1 b_1 + a_2 b_2 + a_3 b_3$$

in terms of coordinates

Given the vector space of SQUARE INTEGRABLE functions with domain (α, β)

and codomain \mathbb{R} , the map

$$(f, g) \mapsto \int_{\alpha}^{\beta} fg$$

specifies an inner product.

A mapping $\cdot : V \times V \rightarrow \mathbb{C}$, where V is a complex vector space, is called a complex inner product if for all $\underline{a}, \underline{b}, \underline{c} \in V$ and $\lambda, \mu \in \mathbb{C}$:

- 1 $\underline{a} \cdot \underline{a} = 0 \Leftrightarrow \underline{a} = \underline{0}$;
- 2 $\underline{a} \cdot \underline{a} \geq 0$;
- 3 $\underline{a} \cdot \underline{b} = \overline{\underline{b} \cdot \underline{a}}$;
- 4 $\underline{a} \cdot (\lambda \underline{b} + \mu \underline{c}) = \lambda \underline{a} \cdot \underline{b} + \mu \underline{a} \cdot \underline{c}$.

Schwarz's Inequality states that, given a real inner product space V ,

$$\underline{x} \cdot \underline{y} \leq (\underline{x} \cdot \underline{x})^{\frac{1}{2}} (\underline{y} \cdot \underline{y})^{\frac{1}{2}}$$

for all $\underline{x}, \underline{y} \in V$. For the space of square integrable functions with the inner product given previously, this becomes

$$\int_{\alpha}^{\beta} fg \leq \left\{ \int_{\alpha}^{\beta} f^2 \right\}^{\frac{1}{2}} \left\{ \int_{\alpha}^{\beta} g^2 \right\}^{\frac{1}{2}}$$

A mapping $\| \cdot \| : V \rightarrow \mathbb{R}$, where V is a real vector space, is called a norm if for all $\underline{a}, \underline{b} \in V$ and $\lambda \in \mathbb{R}$:

- 1 $\| \underline{a} \| = 0 \Leftrightarrow \underline{a} = \underline{0}$;
- 2 $\| \underline{a} \| \geq 0$;
- 3 $\| \lambda \underline{a} \| = |\lambda| \| \underline{a} \|$;
- 4 $\| \underline{a} + \underline{b} \| \leq \| \underline{a} \| + \| \underline{b} \|$ (Triangle Inequality).

In an inner product space the specification

$$\| \underline{a} \| = (\underline{a} \cdot \underline{a})^{\frac{1}{2}}$$

yields a norm satisfying the axioms above. The norm of a geometric vector, given in this way, is just the length of an arrow representing it; by convention, if \underline{a} represents a geometric vector (or a vector field) then its length is denoted by $|\underline{a}|$ or just a .

In G^3 (but not in G^2) we define the vector product $\times : G^3 \times G^3 \rightarrow G^3$ by

$$\underline{OA} \times \underline{OB} = \underline{OA} \cdot \underline{OB} \sin \theta \underline{n},$$

where θ is the angle ($0 < \theta < \pi$) between \underline{OA} and \underline{OB} , and \underline{n} is the *unit vector* (i.e. vector whose norm is 1) orthogonal to \underline{OA} and \underline{OB} , such that \underline{OA} , \underline{OB} and \underline{n} form a right-handed triad

The vector product has the following properties:

- 1 $\underline{i} \times \underline{j} = \underline{k}$, $\underline{j} \times \underline{k} = \underline{i}$, $\underline{k} \times \underline{i} = \underline{j}$, $\underline{i} \times \underline{i} = \underline{j} \times \underline{j} = \underline{k} \times \underline{k} = 0$;
- 2 $\underline{a} \times \underline{b} = -\underline{b} \times \underline{a}$;
- 3 $\underline{a} \times (\underline{b} + \underline{c}) = \underline{a} \times \underline{b} + \underline{a} \times \underline{c}$;
- 4 $\underline{a} \times \underline{b} = (a_2b_3 - a_3b_2)\underline{i} + (a_3b_1 - a_1b_3)\underline{j} + (a_1b_2 - a_2b_1)\underline{k}$;
- 5 $(\underline{a} \times \underline{b}) \times \underline{c} \neq \underline{a} \times (\underline{b} \times \underline{c})$, in general.

$$6 \quad \underline{a} \cdot (\underline{b} \times \underline{c}) = \underline{b} \cdot (\underline{c} \times \underline{a}) = \underline{c} \cdot (\underline{a} \times \underline{b}) = \begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix}$$

$$7 \quad \underline{a} \times (\underline{b} \times \underline{c}) = (\underline{a} \cdot \underline{c})\underline{b} - (\underline{a} \cdot \underline{b})\underline{c}$$

A point in space with Cartesian coordinates (x,y,z) may also be specified by its spherical polar coordinates (r,θ,ϕ) or by its cylindrical polar coordinates (ρ,ϕ,z) . These are related to each other as follows.

$$r^2 = x^2 + y^2 + z^2 \qquad \rho^2 = x^2 + y^2$$

$$x = r \sin \theta \cos \phi = \rho \cos \phi$$

$$y = r \sin \theta \sin \phi = \rho \sin \phi$$

$$z = r \cos \theta$$

In evaluating a VOLUME INTEGRAL the volume element dV is replaced by

$$dx \, dy \, dz \qquad \text{in Cartesian coordinates}$$

$$r^2 \sin \theta \, dr \, d\theta \, d\phi \qquad \text{in spherical polar coordinates}$$

$$\rho \, d\rho \, d\phi \, dz \qquad \text{in cylindrical polar coordinates}$$

The slope of a SCALAR FIELD Ψ is given by the VECTOR FIELD

$$\begin{aligned}\underline{\text{grad}} \Psi &= \frac{\partial \Psi}{\partial x} \underline{i} + \frac{\partial \Psi}{\partial y} \underline{j} + \frac{\partial \Psi}{\partial z} \underline{k} \\ &= \frac{\partial \Psi}{\partial r} \underline{e}_r + \frac{1}{r} \frac{\partial \Psi}{\partial \theta} \underline{e}_\theta + \frac{1}{r \sin \theta} \frac{\partial \Psi}{\partial \phi} \underline{e}_\phi \\ &= \frac{\partial \Psi}{\partial \rho} \underline{e}_\rho + \frac{1}{\rho} \frac{\partial \Psi}{\partial \phi} \underline{e}_\phi + \frac{\partial \Psi}{\partial z} \underline{k},\end{aligned}$$

where \underline{e}_α denotes the vector field of unit norm in the direction of increasing α .

If \underline{v} is a VECTOR FIELD given by

$$\begin{aligned}\underline{v} &= v_x \underline{i} + v_y \underline{j} + v_z \underline{k} \\ &= v_r \underline{e}_r + v_\theta \underline{e}_\theta + v_\phi \underline{e}_\phi \\ &= v_\rho \underline{e}_\rho + v_\phi \underline{e}_\phi + v_z \underline{k},\end{aligned}$$

we may define the SCALAR FIELD

$$\begin{aligned}\text{div } \underline{v} &= \frac{\partial v_x}{\partial x} + \frac{\partial v_y}{\partial y} + \frac{\partial v_z}{\partial z} \\ &= \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 v_r) + \frac{1}{r \sin \theta} \frac{\partial}{\partial \theta} (\sin \theta v_\theta) + \frac{1}{r \sin \theta} \frac{\partial v_\phi}{\partial \phi} \\ &= \frac{1}{\rho} \frac{\partial}{\partial \rho} (\rho v_\rho) + \frac{1}{\rho} \frac{\partial v_\phi}{\partial \phi} + \frac{\partial v_z}{\partial z}.\end{aligned}$$

The LAPLACIAN operator is given by

$$\begin{aligned}\nabla^2 \Psi &= \text{div } \underline{\text{grad}} \Psi \\ &= \frac{\partial^2 \Psi}{\partial x^2} + \frac{\partial^2 \Psi}{\partial y^2} + \frac{\partial^2 \Psi}{\partial z^2} \\ &= \frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial \Psi}{\partial r} \right) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial \Psi}{\partial \theta} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 \Psi}{\partial \phi^2} \\ &= \frac{1}{\rho} \frac{\partial}{\partial \rho} \left(\rho \frac{\partial \Psi}{\partial \rho} \right) + \frac{1}{\rho^2} \frac{\partial^2 \Psi}{\partial \phi^2} + \frac{\partial^2 \Psi}{\partial z^2}.\end{aligned}$$

The following rules are useful:

$$\underline{\text{grad}}(\Psi\Phi) = \Psi \underline{\text{grad}} \Phi + \Phi \underline{\text{grad}} \Psi$$

$$\text{div}(\Psi \underline{v}) = \Psi \text{div} \underline{v} + \underline{v} \cdot \underline{\text{grad}} \Psi$$

$$\text{div}(\Psi \underline{\text{grad}} \Phi) = \Psi \nabla^2 \Phi + \underline{\text{grad}} \Psi \cdot \underline{\text{grad}} \Phi$$

Divergence Theorem (3: pages 14,28)

Let D be a CONVEX DOMAIN in R^2 (or R^3) bounded by the closed CURVE (or closed surface) C. If the coordinates of the VECTOR FIELD \underline{v} are continuous on $D \cup C$ and their first partial derivatives are continuous in D then

$$\int_D \text{div} \underline{v} \, dA = \oint_C \underline{v} \cdot \underline{n} \, ds \quad (D \subset R^2),$$

$$\int_D \text{div} \underline{v} \, dV = \int_C \underline{v} \cdot \underline{n} \, dS \quad (D \subset R^3),$$

where \underline{n} is the unit outward normal to C.

Green's Theorem (W: page 53)

Let D be a CONVEX DOMAIN in R^2 (or R^3) bounded by the closed CURVE (or closed surface) C. If the SCALAR FIELD u has continuous first partial derivatives on $D \cup C$ and continuous second partial derivatives in D then

$$\int_D u \nabla^2 u \, dA = \oint_C u \frac{\partial u}{\partial n} \, ds - \int_D |\underline{\text{grad}} u|^2 \, dA \quad (D \subset R^2),$$

$$\int_D u \nabla^2 u \, dV = \int_C u \frac{\partial u}{\partial n} \, dS - \int_D |\underline{\text{grad}} u|^2 \, dV \quad (D \subset R^3),$$

where $\partial u / \partial n$ is the outward normal derivative of u on C.

3 USEFUL FORMULAS AND THEOREMS

Trigonometric Functions

$$\sin\left(\frac{\pi}{2} - \theta\right) = \cos\theta$$

$$\cos\left(\frac{\pi}{2} - \theta\right) = \sin\theta$$

$$\sin^2\theta + \cos^2\theta = 1$$

$$\sin 2\theta = 2\sin\theta\cos\theta$$

$$\cos 2\theta = \cos^2\theta - \sin^2\theta$$

$$= 2\cos^2\theta - 1 = 1 - 2\sin^2\theta$$

$$\sin 3\theta = 3\sin\theta - 4\sin^3\theta$$

$$\cos 3\theta = 4\cos^3\theta - 3\cos\theta$$

$$\tan 2\theta = \frac{2 \tan\theta}{1 - \tan^2\theta}$$

$$1 + \tan^2\theta = \sec^2\theta$$

$$1 + \cot^2\theta = \operatorname{cosec}^2\theta$$

$$\sin(\alpha \pm \beta) = \sin\alpha\cos\beta \pm \cos\alpha\sin\beta$$

$$\cos(\alpha \pm \beta) = \cos\alpha\cos\beta \mp \sin\alpha\sin\beta$$

$$\tan(\alpha \pm \beta) = \frac{\tan\alpha \pm \tan\beta}{1 \mp \tan\alpha\tan\beta}$$

$$\sin\alpha + \sin\beta = 2\sin \frac{1}{2}(\alpha+\beta)\cos \frac{1}{2}(\alpha-\beta)$$

$$\sin\alpha - \sin\beta = 2\cos \frac{1}{2}(\alpha+\beta)\sin \frac{1}{2}(\alpha-\beta)$$

$$\cos\alpha + \cos\beta = 2\cos \frac{1}{2}(\alpha+\beta)\cos \frac{1}{2}(\alpha-\beta)$$

$$\cos\alpha - \cos\beta = -2\sin \frac{1}{2}(\alpha+\beta)\sin \frac{1}{2}(\alpha-\beta)$$

$$\cot\theta + \tan\theta = 2\operatorname{cosec}2\theta$$

$$\cot\theta - \tan\theta = 2\cot 2\theta$$

$$e^{i\theta} = \cos\theta + i\sin\theta \quad (\text{Euler's formula})$$

Hyperbolic Functions

$$\sinh x = \frac{1}{2}(e^x - e^{-x}) = -i \sin ix$$

$$\cosh x = \frac{1}{2}(e^x + e^{-x}) = \cos ix$$

$$\tanh x = \frac{\sinh x}{\cosh x}$$

$$\cosh^2 x - \sinh^2 x = 1$$

Trigonometric Fourier Series

The trigonometric Fourier series of the function f with domain $[-L, L]$ is given by

$$f(x) \sim \frac{1}{2}a_0 + \sum_{n=1}^{\infty} \left(a_n \cos \frac{n\pi x}{L} + b_n \sin \frac{n\pi x}{L} \right),$$

where

$$a_n = \frac{1}{L} \int_{-L}^L f(x) \cos \frac{n\pi x}{L} dx$$

$$b_n = \frac{1}{L} \int_{-L}^L f(x) \sin \frac{n\pi x}{L} dx$$

Taylor's Theorem (5: page 8, 8: page 10)

If u is a function of one variable which is $n+1$ times continuously differentiable on the interval $[x, x+h]$, then there is a $\theta \in (0, 1)$ such that

$$u(x+h) = u(x) + hu'(x) + \dots + \frac{h^n}{n!} u^{(n)}(x) + \frac{h^{n+1}}{(n+1)!} u^{(n+1)}(x+\theta h).$$

(for $h < 0$, $[x, x+h]$ is replaced by $[x+h, x]$.) If, in addition, $u^{(n+1)}$ is bounded on $[x, x+h]$, i.e.,

$$|u^{(n+1)}(\bar{x})| \leq B \quad \bar{x} \in [x, x+h]$$

for some B , then the remainder term is of order h^{n+1} :

$$\frac{h^{n+1}}{(n+1)!} u^{(n+1)}(x+\theta h) = O(h^{n+1}) \text{ as } h \rightarrow 0.$$

Intermediate Value Theorem (8: page 10)

If a function f is differentiable on the interval $[a,b]$, then for any real number y between $f'(a)$ and $f'(b)$, there is at least one point $x_0 \in [a,b]$ such that $f'(x_0) = y$.

Principle of Superposition (W: page 33)

The solution of the linear equation

$$L[u] = F$$

subject to the linear conditions

$$L_i[u] = f_i \quad i = 1, 2, \dots, n$$

is given by

$$u = u_0 + u_1 + \dots + u_n$$

where u_0 satisfies the problem

$$L[u_0] = F$$

$$L_i[u_0] = 0 \quad i = 1, \dots, n$$

and (for $k = 1, \dots, n$) u_k satisfies the problem

$$L[u_k] = 0$$

$$L_i[u_k] = f_i \delta_{ik}$$

(δ_{ik} is the KRONECKER DELTA).

Extremum Principles for Flux (3: pages 22,23)

Let the SCALAR FIELD w satisfy $\nabla^2 w = -k$ in a domain D , with $w = 0$ on C (the closed boundary of D). Then if w^* is any nonzero SCALAR FIELD which is differentiable on D and continuous on $D \cup C$ such that $w^* = 0$ on C , and \underline{v}^* is any VECTOR FIELD such that $\text{div } \underline{v}^* = -k$ in D , then

$$\frac{k \left(\int_D w^* dA \right)^2}{\int_D |\text{grad } w^*|^2 dA} \leq \int_D w dA \leq \frac{1}{k} \int_D v^{*2} dA.$$

Maximum Principle for Poisson's Equation (W: pages 55,56)

If $\nabla^2 u = -F$ in D , and $F \leq 0$, then

$$\max_{\underline{x} \in D} u(\underline{x}) \leq \max_{\underline{x} \in C} u(\underline{x})$$

where C is the boundary of D .

Finite-Difference Formulas

$$u'(x) \approx \frac{1}{h} [u(x) - u(x-h)] \quad \text{backward-difference formula}$$

$$u'(x) \approx \frac{1}{2h} [u(x+h) - u(x-h)] \quad \text{central-difference formula}$$

$$u'(x) \approx \frac{1}{h} [u(x+h) - u(x)] \quad \text{forward-difference formula}$$

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} \approx \frac{\delta_x^2 u}{(\Delta x)^2} + \frac{\delta_y^2 u}{(\Delta y)^2} \quad \text{five-point formula}$$

(δ_x is the CENTRAL-DIFFERENCE OPERATOR)

$$\begin{aligned} \delta_x^2 u(x) &= u(x+h) - 2u(x) + u(x-h) \\ &= h^2 u''(x) + \frac{1}{12} h^4 u^{(4)}(x) + O(h^6). \end{aligned}$$

The DIFFUSION OPERATOR

$$\frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2} \quad u$$

may be replaced by:

the Crank-Nicolson scheme

$$u_{i,j+1} - u_{i,j} = \frac{\tau}{2} [\delta_x^2 u_{i,j+1} + \delta_x^2 u_{i,j}],$$

the Du Fort and Frankel scheme

$$u_{i,j+1} - u_{i,j-1} = 2\tau [u_{i+1,j} - u_{i,j+1} - u_{i,j-1} + u_{i-1,j}],$$

or the explicit scheme

$$u_{i,j+1} - u_{i,j} = r \delta_x^2 u_{i,j}.$$

Here $r = k/h^2$ is the MESH RATIO, and $u_{i,j}$ denotes $u(ih,jk)$.

Courant-Friedrichs-Lewy (C.F.L.) Condition (8: page 14)

A necessary and sufficient condition for the convergence of a FINITE-DIFFERENCE SCHEME for a HYPERBOLIC equation is that the NUMERICAL DOMAIN OF DEPENDENCE of a point must include the DOMAIN OF DEPENDENCE of the same point.

Lax's Theorem (8: page 16)

Given a PROPERLY POSED, LINEAR, INITIAL VALUE PROBLEM and a linear FINITE-DIFFERENCE REPLACEMENT which is COMPATIBLE with it, the finite-difference scheme is CONVERGENT if it is STABLE.

A linear FINITE-DIFFERENCE REPLACEMENT of an INITIAL VALUE PROBLEM is STABLE if

- (i) the SPECTRAL RADIUS of the matrix corresponding to the finite-difference scheme is not greater than 1 (Matrix Method)

or

- (ii) the solutions to the scheme which have the form $e^{i\beta p h} \xi^q$ satisfy $|\xi| \leq 1$ (von Neumann's method).

Newton's Method

The nonlinear system of equations

$$\underline{f}(\underline{x}) = \underline{0},$$

where $\underline{f}(\underline{x})$ denotes $(f_1(\underline{x}), f_2(\underline{x}), \dots, f_N(\underline{x}))$, may be solved by the linear
ITERATIVE SCHEME

$$J_{\underline{x}^{(n)}} (\underline{x}^{(n)} - \underline{x}^{(n+1)}) = \underline{f}(\underline{x}^{(n)})$$

where $J_{\underline{x}^{(n)}}$ denotes the matrix whose determinant is the JACOBIAN of the functions $\{f_i\}$ at $\underline{x}^{(n)}$.

Chain Rule

If u, x_1, x_2, \dots, x_N are variables which depend on t , and

$$u = f(x_1, \dots, x_N)$$

then

$$\frac{du}{dt} = \sum_{i=1}^N \frac{\partial f}{\partial x_i} \frac{dx_i}{dt}.$$

Fourier Series

Let $\{\phi_n\}$ be an ORTHOGONAL set and f SQUARE INTEGRABLE on $[\alpha, \beta]$ with respect to the weight function ρ , and let the Fourier series for f be given by

$$f \sim \sum_{n=1}^{\infty} c_n \phi_n.$$

Then $\sum_{n=1}^{\infty} c_n^2 \int_{\alpha}^{\beta} \phi_n^2 \rho$ converges and satisfies Bessel's Inequality

$$\sum_{n=1}^{\infty} c_n^2 \int_{\alpha}^{\beta} \phi_n^2 \rho \leq \int_{\alpha}^{\beta} f^2 \rho.$$

Equality holds if and only if the Fourier series for f CONVERGES IN THE MEAN TO f , in which case we have Parseval's Equation

$$\sum_{n=1}^{\infty} c_n^2 \int_{\alpha}^{\beta} \phi_n^2 \rho = \int_{\alpha}^{\beta} f^2 \rho.$$

If $\{\phi_n\}$ is COMPLETE, and

$$f^* \sim \sum_{n=1}^{\infty} c_n^* \phi_n,$$

then

$$\sum_{n=1}^{\infty} c_n c_n^* \int_{\alpha}^{\beta} \phi_n^2 \rho = \int_{\alpha}^{\beta} f f^* \rho.$$

In the following, let L be the operator defined by

$$L : u \mapsto (pu')' - qu,$$

and let u_k denote the k th EIGENFUNCTION of

$$Lu + \lambda \rho u = 0 \text{ in } (\alpha, \beta)$$

$$u(\alpha) = u(\beta) = 0.$$

Lagrange's Identity

$$\int (uLv - vLu) = [p(uv' - vu')]$$

over any interval in which u and v are twice CONTINUOUSLY DIFFERENTIABLE.

Theorem A

The set $\{u_k\}$ is COMPLETE for the space of functions SQUARE INTEGRABLE on (α, β) with weight function ρ .

Theorem B

If f is continuous on $[\alpha, \beta]$, $f(\alpha) = f(\beta) = 0$ and

$$\int_{\alpha}^{\beta} (pf'^2 + qf^2)$$

exists, the Fourier series of f in terms of $\{u_k\}$ is UNIFORMLY CONVERGENT to f on $[\alpha, \beta]$.

Oscillation Theorem

The eigenfunction u_k has precisely $k-1$ zeros in the open interval (α, β) , and the zeros of successive eigenfunctions interlace.

Tests for Uniform ConvergenceCAUCHY'S TEST

A necessary and sufficient condition for a sequence of functions u_n with domain I to be uniformly convergent is that given any $\varepsilon > 0$ there exists an integer N_ε independent of $x \in I$ such that

$$|u_n(x) - u_m(x)| < \varepsilon$$

for all $n, m > N_\varepsilon$ and all $x \in I$.

WEIERSTRASS'S M-TEST

This is a sufficient though not necessary condition. If each term of the series

$$f(x) = \sum_{n=1}^{\infty} u_n(x) \quad x \in I$$

is positive and

$$u_n(x) < M_n \quad \forall x \in I, n \in \mathbb{Z}^+,$$

where each M_n is independent of x , and if

$$\sum_{n=1}^{\infty} M_n$$

is convergent, then the given series for f is uniformly convergent.

Bessel Functions

J_m is a solution of BESSEL'S EQUATION OF ORDER m , and is bounded at the origin.

$$J_m(t) = \sum_{k=0}^{\infty} \frac{(-1)^k (\frac{1}{2}t)^{m+2k}}{k! (m+k)!}$$

$$\frac{d}{dt} [t^{-m} J_m(t)] = -t^{-m} J_{m+1}(t)$$

$$\frac{d}{dt} [t^m J_m(t)] = t^m J_{m-1}(t)$$

$$\int_0^1 x [J_m(\sqrt{\lambda_k^{(m)}} x)]^2 dx = \frac{1}{2} [J_{m+1}(\sqrt{\lambda_k^{(m)}})]^2 \text{ where } J_m(\sqrt{\lambda_k^{(m)}}) = 0$$

$$J'_m(t) = \frac{1}{2} [J_{m-1}(t) - J_{m+1}(t)]$$

$$\frac{2t}{m} J_m(t) = J_{m+1}(t) + J_{m-1}(t)$$

4 TABLE OF INDEFINITE INTEGRALS

Function	Indefinite Integral with respect to x
$(x-a)^n$	$\frac{(x-a)^{n+1}}{n+1} \quad (n \neq -1)$
$\frac{1}{x-a}$	$\ln x-a $
$\frac{1}{x^2+a^2}$	$\frac{1}{a} \tan^{-1} \frac{x}{a}$
$\frac{1}{a^2-x^2}$	$\frac{1}{2 a } \ln \left \frac{a+x}{a-x} \right $
$\frac{2x+a}{x^2+ax+b}$	$\ln x^2+ax+b $
$\frac{2x+a}{(x^2+ax+b)^n}$	$\frac{(x^2+ax+b)^{1-n}}{1-n} \quad (n \neq 1)$
$\frac{1}{\sqrt{(a^2-x^2)}}$	$\sin^{-1} \frac{x}{ a }$
$\frac{1}{\sqrt{(a^2+x^2)}}$	$\sinh^{-1} \frac{x}{ a }$
$\frac{1}{\sqrt{(x^2-a^2)}}$	$\pm \cosh^{-1} \left \frac{x}{a} \right \quad (\text{sign that of } x)$
$\sin x$	$-\cos x$
$\cos x$	$\sin x$
$\tan x$	$\ln \sec x $
$\cot x$	$\ln \sin x $
$\sec x$	$\ln \sec x + \tan x $
$\operatorname{cosec} x$	$\ln \tan \frac{1}{2}x $
$\sec^2 x$	$\tan x$
$\operatorname{cosec}^2 x$	$-\cot x$
e^{ax}	$\frac{1}{a} e^{ax}$
a^x	$\frac{a^x}{\ln a}$

$\sinh x$	$\cosh x$
$\cosh x$	$\sinh x$
$\tanh x$	$\ln(\cosh x)$
$\coth x$	$\ln \sinh x $
$\operatorname{sech} x$	$\frac{1}{2} \tan^{-1}(e^x)$
$\operatorname{cosech} x$	$\ln \tanh \frac{1}{2}x $
$\operatorname{sech}^2 x$	$\tanh x$
$\operatorname{cosech}^2 x$	$-\coth x$
$\sin rx \cos nx \ (r^2 \neq n^2)$	$-\frac{\cos(r-n)x}{2(r-n)} - \frac{\cos(r+n)x}{2(r+n)}$
$\sin rx \sin nx \ (r^2 \neq n^2)$	$\frac{\sin(r-n)x}{2(r-n)} - \frac{\sin(r+n)x}{2(r+n)}$
$\cos rx \cos nx \ (r^2 \neq n^2)$	$\frac{\sin(r-n)x}{2(r-n)} + \frac{\sin(r+n)x}{2(r+n)}$
$e^{ax} \cos(bx + c)$	$\frac{e^{ax}}{a^2 + b^2} [\operatorname{acos}(bx+c) + b \sin(bx+c)]$
$e^{ax} \sin(bx + c)$	$\frac{e^{ax}}{a^2 + b^2} [a \sin(bx+c) - b \cos(bx+c)]$

5 TABLE OF DEFINITE INTEGRALS

In the following m and n are integers.

$$\int_0^{\pi} \sin mx \sin nx \, dx = 0$$

$$m \neq n$$

$$\int_0^{\pi} \sin nx \, dx = \frac{\pi}{2}$$

$$n \neq 0$$

$$\int_0^{\pi} \cos mx \cos nx \, dx = \begin{cases} 0 \\ \pi \\ \frac{1}{2}\pi \end{cases}$$

$$m \neq n$$

$$m = n = 0$$

$$m = n \neq 0$$

$$\int_{-\pi}^{\pi} \cos mx \cos nx \, dx = \begin{cases} 0 \\ \pi \end{cases}$$

$$m \neq n$$

$$m = n \neq 0$$

$$\int_{-\pi}^{\pi} \sin mx \cos nx \, dx = 0$$

$$\int_{-\pi}^{\pi} \cos nx \, dx = \begin{cases} 0 \\ 2\pi \end{cases}$$

$$n \neq 0$$

$$n = 0$$

$$\int_{-\pi}^{\pi} \sin nx \, dx = 0$$

$$\int_{-\infty}^{\infty} e^{-x^2} \, dx = \sqrt{\pi}$$